

NOTES

Emergence in Italy of a *Neisseria meningitidis* Clone with Decreased Susceptibility to Penicillin

Paola Stefanelli, Cecilia Fazio, Arianna Neri, Tonino Sofia, and Paola Mastrantonio*

Department of Infectious, Parasitic and Immuno-Mediated Diseases, Istituto Superiore di Sanita', Rome, Italy

Received 8 March 2004/Returned for modification 6 April 2004/Accepted 13 April 2004

A rise in invasive diseases due to *Neisseria meningitidis* C:2b:P1.5 with decreased penicillin susceptibility occurred in Italy during the last 2 years. Real-time PCR identified the Penⁱ phenotype, and the *penA* sequence revealed the mosaicism of the gene. Molecular analyses assigned the isolates to a single emergent clone of the hypervirulent A4 cluster.

In Italy, the incidence of meningococcal disease has been consistently low (about 0.3 to 0.4 per 100,000 inhabitants per year). Throughout the 1990s, *Neisseria meningitidis* serogroup C strains represented fewer than 30% of all isolated meningococci. In 2002 to 2003, we observed an increase in serogroup C meningococcal disease, which became responsible for 42.5% of culture-confirmed cases. Interestingly, most of these isolates have the antigenic phenotype C:2b:P1.5 and show intermediate susceptibility to penicillin (Penⁱ) (0.06 µg/ml > MIC < 1 µg/ml), due to sequence changes in the penicillin binding protein 2 *penA* gene. Penⁱ meningococci have been reported and are being monitored in several countries (3, 5, 8). The attention is driven by the concept that strains with decreased susceptibility may still be evolving and in the process of acquiring further alterations in the *penA* gene, anticipating the appearance of resistant strains, as observed for *Streptococcus pneumoniae* and *Neisseria gonorrhoeae* (9, 12). In Italy, before 2002 Penⁱ meningococci accounted for 7.5% of the isolates, but since then the percentage has risen significantly (to 27.4%). All Penⁱ strains showed a mosaic structure in the transpeptidase region of the *penA* gene (1), and a recently validated real-time PCR assay (10) has been used for rapid confirmation of this phenotype. In this study, all C:2b:P1.5 Penⁱ meningococci isolated in Italy were analyzed by *penA* sequencing, multilocus sequence typing, and pulsed-field gel electrophoresis (PFGE) to determine whether the recent increase in the number of cases due to this phenotype correlates with the emergence of a single clone and whether it circulates in other European countries.

A total of 214 invasive *N. meningitidis* strains were received at the reference laboratory of the National Surveillance of Meningococcal Meningitis between January 2002 and December 2003. All isolates were typed (7), and of 214 *N. meningitidis* isolates received, 91 were serogroup C and 38 of these (42.2%) belonged to phenotype C:2b:P1.5. Among the remaining serogroup C strains, 21% had the phenotype C:2a:P1.5, which up to

then was the most frequent phenotype among C strains, and the others showed a variety of different sero- and subtypes.

The recently described real-time PCR (10) was used to discriminate between penicillin-susceptible (Pen^s) and Penⁱ strains. Two hybridization probes were used to distinguish the wild-type *penA* gene in the Pen^s meningococci from the mutated gene at codon 566 in Penⁱ strains. Thermal analysis of probe hybridization revealed melting temperatures of 45.5 and 55°C for the Pen^s and Penⁱ strains, respectively, as shown in Fig. 1A. MICs of penicillin were also assessed by use of the E-test (AB Biodisk) on Mueller-Hinton agar (Oxoid) supplemented with 5% sheep blood, and the majority of C:2b:P1.5 strains (78.4%) showed MICs of ≥0.094 µg/ml, whereas a few serogroup C strains (24%) with other sero- or subtypes were Penⁱ.

The Penⁱ phenotype was finally confirmed by sequencing the *penA* PCR products in the transpeptidase domain of the encoding gene. Analysis of the sequences was performed with the Accelrys Wisconsin Genetics Computer Group package.

The *penA* sequence analysis showed a short DNA region, between nucleotides 1364 and 1545, with 98% identity to the sequence derived from *Neisseria perflava/sicca* (accession number X76422) and 100% homology with all *penA* sequences from Spanish C:2b:P1.5,2 Penⁱ strains deposited in the National Center for Biotechnology Information databank (<http://www.ncbi.nlm.nih.gov>). Conversely, the *penA* genes of Penⁱ meningococci with other serotypes or serosubtypes appear to have acquired larger blocks of DNA, often from more than one commensal *Neisseria* species along all of the transpeptidase domain (Fig. 1B). The replacement of the short region between nucleotides 1364 and 1545 in the C:2b:P1.5 strains seems to be sufficient to produce a form of the protein with a lower affinity to penicillin.

The spread of Penⁱ C:2b:P1.5 meningococci over the country was more remarkable in the first 6 months of 2003, when twice as many were isolated as in the previous year. This sharp increase led us to carry out a molecular analysis of these strains to confirm the circulation of a clone. According to multilocus sequence typing results, obtained by the methodology described by Maiden et al. (6) (<http://neisseria.org/nm/typing>

* Corresponding author. Mailing address: Department of Infectious, Parasitic & Immuno-mediated Diseases, Istituto Superiore di Sanita', Viale Regina Elena 299, 00161-Rome, Italy. Phone: 390649902335. Fax: 390649387112. E-mail: pmastran@iss.it.




FIG. 1. (A) Examples of T_m curves for *Pen^S* and *Pen^I* *N. meningitidis* phenotypes obtained by real-time PCR assay with the mutated 566 codon in the *penA* gene. (B) Mosaic *penA* genes of *Pen^I* *N. meningitidis* strains. Each line indicates the proposed origin of the exogenous DNA blocks in the transpeptidase domain of the gene. The arrowheads show the position of the mutated 566 codon used as a marker of *penA* translocation in the real-time PCR assay. nt, nucleotides; AC#, accession number.

TABLE 1. Characteristics of *N. meningitidis* C:2b:P1.5 strains belonging to the ST8/A4 cluster isolated between January 2002 and December 2003

No. of strains	Yr of isolation	Patient age(s) (yr)	MIC of penicillin ($\mu\text{g/ml}$)	Pulse type
1	2002	1	0.094	PTA
4	2002	3, 4, 17, 19	0.125	PTA
1	2002	18	0.125	PTA1
2	2002	4, 7	0.19	PTA
6	2003	<1, 4, 15, 24, 49, NK ^a	0.094	PTA
1	2003	<1	0.094	PTA2
5	2003	4, 11, 12, 17, 35	0.125	PTA
1	2003	<1	0.125	PTA2
2	2003	23, 46	0.125	PTA1
4	2003	8, 30, 60, 62	0.19	PTA
2	2003	21, NK	0.25	PTA
1	2003	18	0.38	PTA

^a NK, not known.

/mlst/), all of the C:2b:P1.5 Penⁱ meningococci were assigned to the ST8/A4 cluster (Table 1), which is one of the two hyper-virulent lineages responsible for most of serogroup C disease worldwide (the other is ST11). DNA macrorestriction fragments generated with the NheI restriction enzyme and analyzed by PFGE, as previously described (4), showed one main pulse type (PTA) and two subclones with two or three minor differences attributable to point mutations, named PTA1 and PTA2 (Fig. 2). When BglII was used, all of the strains showed the same pulse type (data not shown). One Penⁱ strain with




FIG. 2. PFGE profiles of genomic DNAs from *N. meningitidis* C:2b:P1.5 strains after digestion with endonuclease NheI. The lambda ladder DNA marker (New England Biolabs) (lane MW) was used as a molecular size standard (48.5 kb). Lane 1, pulse type PTA; lane 2, PTA1; lane 3, PTA2. The arrows identify the band differences.

phenotype C:2b:nst, isolated in the first months of 2003, belonged to the same clone. The lack of identification of a sero-subtype in this strain was due to the presence of IS1301 in the *porA* gene and its subsequent inactivation, as demonstrated by sequencing (data not shown).

Interestingly, the pulse type PTA found in most of the examined C:2b:P1.5 strains is identical to fingerprint pattern 2 (PT2) described by Arreaza et al. (2) for C:2b:P1.5,2 epidemic Spanish strains, just as our PTA1 resembles PT1 described in the same paper. Unlike the Spanish PT1 strains, the PTA1 strains are able to cause disease in all age groups and not predominantly in children under 2 years of age. In contrast, the subclone PTA2 is a unique profile found only in two strains, both of which were responsible for meningitis in infants.

All of these findings seem to confirm that the recent increase of meningococcal disease caused by *N. meningitidis* C:2b:P1.5 in Italy is due to the spread of a single emergent clone with decreased penicillin susceptibility and belonging to the hyper-virulent cluster A4. Although the first strain C:2b:P1.5, ST8/A4 cluster appeared in Italy in 1998, in the following 3 years all meningococci with phenotype C:2b:P1.5 belonged to ST1860, a new ST of the ET37 complex detected in Italy (11). None of these strains were Penⁱ, in contrast to most of the serogroup C ET37 strains isolated in other countries. In 2002, Penⁱ meningococci of phenotype C:2b:P1.5, ST8/A4 cluster suddenly reappeared and rapidly spread all over the country.

We speculate that the Spanish clone might have been imported in the second half of the 1990s. In fact, one Penⁱ strain with phenotype C:2b:P1.5,2 ST8/A4 and pulse type PTA1 was isolated in Italy in 1996, and it showed the same molecular characteristics as the Spanish strains. After this episode, for reasons difficult to understand and linked to the evolution and dynamics of the meningococcal population, it took some years to settle in, with a minor modification in the outer membrane proteins encoded by the *porA* gene. However, once established, the phenotype C:2b:P1.5 has become so fit as to spread very rapidly and to successfully compete with the other phenotypes. The acquisition of a very small exogenous DNA fragment might have provided an advantage in terms of fitness compared to the other Penⁱ phenotypes. It is important to underline that there is a direct relationship between the increase in serogroup C strains causing meningococcal disease in Italy and the eight-fold increase of meningococci with decreased susceptibility to penicillin as a result of the spread of this virulent clone. It will be of extreme importance to closely monitor the endemic circulation of this phenotype in order to plan specific vaccination programs in a timely fashion.

We thank the microbiologists of the hospital laboratories participating in the Italian National Surveillance of Bacterial Meningitis for isolating the strains and sending them to the Reference Laboratory of the Istituto Superiore di Sanità (Rome). This study made use of the Neisseria MultiLocus Sequence Typing website (<http://neisseria.mlst.net>) developed by Man-Suen Chan and sited at the University of Oxford.

The development of the Neisseria MultiLocus Sequence Typing website is funded by the Wellcome Trust. This work was partially funded by Ministero della Salute (Italy), Programma per la Ricerca Finalizzata 2002.

REFERENCES

1. Antignac, A., P. Kriz, G. Tzanakaki, J. M. Alonso, and M. K. Taha. 2001. Polymorphism of *Neisseria meningitidis* *penA* gene associated with reduced susceptibility to penicillin. *J. Antimicrob. Chemother.* **47**:285–296.
2. Arreaza, L., S. Berrón, S. Fernández, M. I. Santiago, A. Malvar, and J. A. Vázquez. 2000. Investigation for a more virulent variant among the C:2b:P1.2,5 Spanish meningococcal epidemic strains by molecular epidemiology. *J. Med. Microbiol.* **49**:1079–1084.
3. Berrón, S., and J. A. Vázquez. 1994. Increase in moderate penicillin resistance and serogroup C in meningococcal strains isolated in Spain. Is there any relationship? *Clin. Infect. Dis.* **18**:161–165.
4. Hartstein, A., C. Phelps, and A. Lemonte. 1995. Typing of sequential bacterial isolates by pulsed-field gel electrophoresis. *Diagn. Microbiol. Infect. Dis.* **22**:309–314.
5. Kyaw, M. H., J. C. Bramley, S. Clarke, et al. 2002. Prevalence of moderate penicillin resistant invasive *Neisseria meningitidis* infection in Scotland, 1994–9. *Epidemiol. Infect.* **128**:149–156.
6. Maiden, M. C. J., J. A. Bygraves, E. Feil, et al. 1998. Multilocus sequence typing: a portable approach to the identification of clones within populations of pathogenic microorganisms. *Proc. Natl. Acad. Sci. USA* **95**:3140–3145.
7. Mastrantonio, P., P. Stefanelli, C. Fazio, T. Sofia, A. Neri, G. La Rosa, C. Marianelli, M. Muscillo, M. G. Caporali, and S. Salmasso. 2003. Serotype distribution, antibiotic susceptibility, and genetic relatedness of *Neisseria meningitidis* strains recently isolated in Italy. *Clin. Infect. Dis.* **36**:422–428.
8. Richter, S. S., K. A. Gordon, P. R. Rhomberg, M. A. Pfaller, and R. N. Jones. 2001. *Neisseria meningitidis* with decreased susceptibility to penicillin: report from the SENTRY antimicrobial surveillance program, North America, 1988–99. *Diagn. Microbiol. Infect. Dis.* **41**:83–88.
9. Spratt, B. G., and K. D. Cromie. 1988. Penicillin-binding proteins of gram-negative bacteria. *Rev. Infect. Dis.* **10**:699–711.
10. Stefanelli, P., A. Carattoli, A. Neri, C. Fazio, and P. Mastrantonio. 2003. Prediction of decreased susceptibility to penicillin of *Neisseria meningitidis* strains by real-time PCR. *J. Clin. Microbiol.* **41**:4666–4670.
11. Stefanelli, P., C. Fazio, A. Neri, T. Sofia, and P. Mastrantonio. 2003. First report of capsule replacement among electrophoretic type 37 *Neisseria meningitidis* strains in Italy. *J. Clin. Microbiol.* **41**:5783–5786.
12. Tomasz, A. 1986. Penicillin-binding proteins and the antibacterial effectiveness of beta-lactam antibiotics. *Rev. Infect. Dis.* **8**:S260–S278.